

GREENHOUSE GAS FROM DAIRY MANURE MANAGEMENT AT THE FARMSTEAD

Part 3: DAIRY MANURE MANAGEMENT IMPACT ON NITROUS OXIDE

August 2017

Daily spreading of manure is being reduced since it can result in water quality and nutrient recycling concerns. Long-term manure storage to avoid spreading when the potential for runoff occurs during frozen, snow covered or saturated conditions is encouraged. This helps to maximize nutrient recycling and to reduce nutrient and potential pathogen losses to ground and surface water. Long-term storage is a water quality best management practice. However it can increase a farm's greenhouse gas (GHG) footprint by releasing GHGs— methane (CH₄) with a Global Warming Potential (GWP) of 34 and Nitrous Oxide (N₂O) with a GWP of 298. Since the GWP of GHG are different they are often expressed as carbon dioxide equivalents (CO₂eq). The calculations and justifications used to quantify N₂O emissions from several aerobic (free oxygen present) manure management methods are shown below.

Quantifying Daily Manure-Based N₂O Emissions per Cow

The daily N₂O emissions converted into CO₂eq can be estimated using Equation 1.3 from the IPCC (2006) and EPA (2009) from N₂O emissions from several different manure management methods. Yearly emissions can be estimated by multiplying the average daily emissions by 365 days.

The two variables that can be controlled by farm management are the daily mass of nitrogen (N) excreted by the animals related to their feed diets and the emission factor (EF₃) depending on the manure management system. The N₂O emissions from manure management are less defined than CH₄ emissions with an estimated uncertainty range of a factor of 2 (200%) (IPCC, 2006). N₂O is emitted when the complete biological oxidation of nitrogen to nitrate (NO₃) is interrupted. Table 1.3 provides a summary of the EF₃ values from the IPCC (2006) and EPA (2016) documents for the N₂O emission factor values for various manure management BMPs.

Although the EF₃ values are less than the MCF values used for CH₄ emission N₂O is 8.7 times the GWP of CH₄ so the GWP from aerobic manure management practices can be significant. Of these manure management practices the most aerobic manure management methods have the highest potential for N₂O emission. Processing manure with solid-liquid separation (SLS) to remove a portion of the N from the solid portion sent to the long-term storage would significantly reduce the N₂O emissions from a more aerobic (actively composted) system since the liquid long-term storage would still contain most of the N leaving much less N in the solid storage (Gooch et al., 2005).

Equation 1.3

$$CO_{2eq} = 298 \text{ CO}_2/N_2O \text{ GWP} \times EF_3 \times 44 \text{ N}_2O/28 \text{ N}_2O-N \times N \text{ excreted/cow-day}$$

CO₂eq = Equivalent GWP expressed as carbon dioxide

298 = GWP factor for N₂O

EF₃ = Emission Factor for N₂O emissions from manure management (see Table 1.3)

N Excreted /cow-day = ~0.99 lbs./cow – day (ASAE).

Table 1.3 Approximate emission factors¹ for N₂O (lb. N₂O-N/lb. N)

EF ₃	Manure Management BMP	Condition
0	Daily spread	N/A
0.005	Solid storage	N/A
0.005	Liquid/slurry	With natural crust
0	Liquid/slurry	Without natural crust
0.002	Pit storage	Below animal confinement
0.01	Bedded pack	No mixing
0.07	Bedded pack	Active mixing (composted bedded pack)
0.006	Compost pile	Static
0.01	Compost windrow	Infrequent turning
0.1	Compost windrow	Frequent turning
0	Anaerobic digestion	N/A

¹Source: IPCC (2006) and EPA (2009)

Nitrogen removal rates for various SLS technologies also vary depending on technology and management. See fact sheet 2 for information on CH₄ emissions with these manure management systems. The combination of the GWP of both CH₄ and N₂O emissions need to be included in determining the impact of the manure

management system on GHG emissions. Fact Sheet 4 combines both the CH₄ and N₂O emissions from several common dairy manure management systems. Additional fact sheets discuss ways to reduce GHG emissions although a specific farm may only be able to implement certain ones depending on their existing facilities.

AUTHORS

Peter Wright, PE pew2@cornell.edu (585) 314-5314 *Curt Gooch*, PE cag26@cornell.edu (607) 225-2088

FACT SHEET SERIES: 1 HOW ARE GREENHOUSE GASES GENERATED?, 2 DAIRY MANURE MANAGEMENT IMPACT ON METHANE, 3 DAIRY MANURE MANAGEMENT IMPACT ON NITROUS OXIDE, 4 COMBINING METHANE AND NITROUS OXIDE EMISSIONS FROM DAIRY MANURE MANAGEMENT, 5 GHG REDUCTION FROM CRUSTS ON STORAGE, 6 GHG REDUCTION FROM LIMITING SUMMER STORAGE, 7 GHG FROM SOLID STORAGE SYSTEMS, 8 GHG REDUCTION FROM SOLID/LIQUID SEPARATION, 9 GHG REDUCTION FROM AN IMPERMEABLE COVER, 10 GHG REDUCTION FROM AN ANAEROBIC DIGESTION SYSTEM.

References:

- ASAE D384.2 MAR2005 (R2010) Manure Production and Characteristics ASABE, 2950 Niles Road, St. Joseph, MI 49085-9659, USA
- Environmental Protection Agency. February 4, 2009. Technical Support Document For Manure Management Systems: Proposed Rule for Mandatory Reporting Of Greenhouse Gases, Climate Change Division Office of Atmospheric Programs U.S. Environmental Protection Agency
- Environmental Protection Agency. March 2011. Final Protocol for Quantifying and Reporting the Performance of Anaerobic Digestion Systems for Livestock Manures, Prepared by: Eastern Research Group, Inc. 110 Hartwell Avenue Lexington, MA 02421
- Environmental Protection Agency. April 15, 2016. Inventory of U.S. Greenhouse Gas Emissions and Sinks: 1990 – 2014, <https://www3.epa.gov/climatechange/Downloads/ghgemissions/US-GHG-Inventory-2016-Main-Text.pdf>
- Gooch, C.A., S.F. Inglis, and K.J. Czymmek. 2005. Mechanical Solid-Liquid Manure Separation: Performance Evaluation on Four New York State Dairy Farms – A Preliminary Report. Presented at the 2005 ASAE Annual International Meeting, July 17 – 20, 2005. ASAE Paper No. 05-4104. ASAE 2950 Niles Road, St. Joseph, MI 49085-9659.
- Intergovernmental Panel on Climate Change (IPCC) Tier 2 method from the 2006 IPCC Guidelines for National GHG Inventories, Volume 4, Chapter 10:

This material is based upon work that is supported by the National Institute of Food and Agriculture, U.S. Department of Agriculture, under award number 2013-68002-20525